

Original Article: Carvacrol Alleviated Negative Effects of Diabetes on Inflammation and Oxidation by Modulation in Gene Expression of Inflammatory and Antioxidant System in Diabetic Rat Model



Seyed Ashkan Tabibzadeh Dezfuli¹, Mohanna Ehsani², Omid Lakzaei Azar^{3*}

1. Assistant Professor, Trauma and Emergency Medicine Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

2. Department of Microbiology, Faculty of Basic Sciences, Lahijan Branch, Islamic Azad University, Lahijan, Iran.

3. Young Researchers and Elite Club, Lahijan Branch, Islamic Azad University, Lahijan, Iran.



Citation Tabibzadeh Dezfuli SA, Ehsani M, Lakzaei Azar O. Carvacrol Alleviated Negative Effects of Diabetes on Inflammation and Oxidation by Modulation in Gene Expression of Inflammatory and Antioxidant System in Diabetic Rat Model. GMJ Medicine. 2017; 1(1):15-20.

doi <https://doi.org/10.22034/GMJM.2017.01.15>



Article info:

Received: 25 July 2017

Accepted: 11 October 2017

Available Online: 25 September 2017

Checked for Plagiarism: Yes

Peer reviewers approved by:

Dr. Melika Andrew

Editor who approved publication:

Prof. Dr. Nanuli Doreulee

Keywords:

Antioxidant enzymes,
Diabetes, Pro-inflammatory
cytokines

ABSTRACT

Background: Diabetes has been known as a prevalence disorder and the use of common drugs has been faced many issues with multiple limitations. This study aimed to evaluate the use of carvacrol, as a novel agent, for treatment of diabetes.

Materials and Methods: A single dose of streptozotocin (55 mg/kg body weight) was used to induce the diabetes in rats. The animals were grouped into five groups including 1. Control healthy animals; 2. Diabetic controls; 3, 4 & 5. Diabetic animals given carvacrol (5, 10 and 15 mg/kg body weight/day) in neutral sterile olive oil solution oral gavage, respectively. The levels of Malondialdehyde (MDA) and catalase, Superoxide Dismutase (SOD) and Glutathione Peroxidase (GPX) activities were evaluated. The levels of IL-1 β , IL-6 and TNF- α expression in liver were assessed.

Results: Administration of streptozotocin increased levels of MDA, IL-1 β , IL-6 and TNF- α and also decreased activities of catalase, SOD and GPX (P<0.05). Oral administration of carvacrol, especially 15 mg/kg body weight/day, could decrease levels of MDA, IL-1 β , IL-6 and TNF- α and also increase activities of catalase, SOD and GPX in comparison to diabetic control (P<0.05).

Conclusion: Carvacrol could decrease or alleviate the negative effects of carvacrol on inflammation and antioxidant status that could be attributed to antioxidant properties. It could be recommended to apply carvacrol in commercial prescription in combination with other agents or as a single agent for treatment of diabetes.

* Corresponding Author:

Omid Lakzaei Azar, PhD

Address: Young Researchers and Elite Club, Lahijan Branch, Islamic Azad University, Lahijan, Iran.

E-mail: omid.lakzaeazar@yahoo.com

Introduction

Diabetes mellitus is a complex disease because it is a metabolic disorder that is involved with oxidation and inflammation [1]. It has been accepted that inflammation has roles in type 1 and type 2 diabetes mellitus [2, 3]. Studies have shown increased inflammatory markers in patients with diabetes [4]. Diabetic nephropathy has been known as one of the most common complications in diabetes and it is also main reason for end-stage renal disease in all over world [5]. Glucose is entered into cells in independent form of insulin and accumulated glucose increases inflammation, cell oxidation and apoptosis [6].

Pathogenesis of diabetic nephropathy could be due to binary of hemodynamic changes and excessive hyperglycemia, resulting in increased inflammation and oxidative stress [7, 8]. Increased oxidative stress and formation of pro-inflammatory cytokines happens during diabetes [9]. During Inflammation, inflammatory reactions raise levels of Tumor Necrosis Factor- α (TNF- α), Interlukin-1 β (IL-1 β) and IL-6 that interact with proteins [10].

The TNF- α has been known as an inflammatory cytokine that initiates inflammation. The IL-1 β is a pro-inflammatory cytokine that calls of neutrophils into the region of infection [11]. On the other hand, oxidative stress is known to have excessive importance in diabetes. There is an imbalance between the formation and neutralization of Reactive Oxygen Species (ROS) including highly reactive hydroxyl radicals' superoxide anion, peroxy radicals, singlet oxygen, peroxy nitrite, and hydrogen peroxide. Enzymatic and non-enzymatic antioxidants help to treat and/or prevent the diabetes and its related complications [12].

Synthetic drugs have also been used to treat the diabetes, but its use has faced with limitations due to side effects. Plant derivations have also been used to treat the different diseases. Carvacrol, one monoterpene phenol, that is extensively existed in some species including *Origanum*, *Satureja*, *Thymbra*, *Thymus*, and *Corydolithymus* [13]. Carvacrol has been reported to have some pharmacological features including antioxidant [14], anti-inflammatory [15], antitumor [16], and antimicrobial [17] activity. It seems that carvacrol could improve diabetes due to its antioxidant and anti-inflammatory. We aim to introduce the carvacrol as a novel agent for treatment of diabetes. This study was thus conducted to

evaluate the effects of carvacrol on inflammatory and antioxidant responses of diabetic rats.

Materials and Methods

Materials

Carvacrol was prepared from Fluka, Chemika, Sigma-Aldrich (St Quentin Fallavier, France) in purity of 95%. Streptozotocin was also prepared from Sigma-Aldrich Company. All other chemical agents used were analytical grade and prepared from standard commercial suppliers.

Animals

Adult male Wistar rats (220 \pm 20 g) were from the Pasture Institute, Tehran-Iran. Animals were kept in lighting program of 12 h/12 h light/dark cycle. Animals had ad libitum access to food and water. All the used principles were approved by the Guide for the Care and Use of Laboratory Animals, USA, 1986.

Induction of diabetes

A single dose of streptozotocin (55 mg/kg body weight) in 0.1 M cold citrate buffer (pH 4.5) was intra peritoneal administered to fasted rats. Control animals received citrate buffer alone. In overnight, animals consumed 5% glucose solution in order to overcome on hypoglycemia. Seventy-two hours after administration of streptozotocin, glucose was assessed by glucometer and rats with glucose concentrations >300 mg/dL were used as diabetic and studied. Animals were grouped one week after administration of streptozotocin and lasted for 7 days.

Experimental design

The animals (n=50) were grouped into five treatments, each comprising of ten animals, including: Group 1. Control animals giving 0.1M citrate buffer (pH=4.5); Group 2. Diabetic controls; Group 3. Diabetic animals given carvacrol (5 mg/kg Body weight/day) in neutral sterile olive oil solution oral gavage; Group 4. Diabetic animals given carvacrol (10 mg/kg Body weight/day) in neutral sterile olive oil solution oral gavage; Group 5. Diabetic animals given carvacrol (15 mg/kg Body weight/day) in neutral sterile olive oil solution oral gavage.

Evaluation of blood glucose

In day 7 of trial, the Wistar rats were fasted at overnight, anaesthetized and sacrificed, by cervical dislo-

cation. The blood samples was gathered in tubes with EDTA for investigation of plasma glucose.


Assessment of antioxidant activity

After killing rats, the livers were removed and washed with isotonic saline. The liver tissue was homogenated and used in 5% (w/v) potassium phosphate buffer (0.1 M, pH 7.4) by a homogenizer. The homogenate sample was then centrifuged in 16000×g for 20 min in order to remove the nuclei and cell debris. The supernatant was applied for measurement of lipid peroxidation and antioxidant activity as previously reported [18]. In order to evaluate the Malondialdehyde (MDA), TBARS contents of the samples were measured from a standard curve by 1,1,3,3-tetramethoxypropane.

Catalase activity was measured by the molar extinction coefficient of 43.6 M⁻¹ cm⁻¹ for H₂O₂ and reported as μmol H₂O₂ consumed/min per milligram of protein. To evaluate the Superoxide Dismutase (SOD) and Glutathione Peroxidase (GPx) activities, diagnostic kits of RAN-SOD and ANSEL (RANDOX, UK) were used and the values were reported as unit/mg protein. Protein level of the samples was assessed as reported by Bradford [19].

Real-time RT-PCR for IL-1β, IL-6 and TNF-α

Parts of liver samples were used to evaluate the Real-time RT-PCR for IL-1β, IL-6 and TNF-α. It was investigated as reported by Kha et al. [20]. The primers sequences were IL-1β, forward (5'-CACCTTCTTTTCCTTCATCTTTG-3') and reverse (5'-GTCGTTGCTTGTCTCTCCTTGTA-3'), IL-6, forward (5'-TGATGGATGCTTCCAAACTG-3') and reverse (5'-GAGCATTGGAAGTTGGGGTA-3'), TNF-α, forward (5'-ACTGAACTTCGGGGTGATTG-3') and reverse (5'-GCTTGGTGGTTTGCTACGAC-3') and GADPH forward (5'-GTATTGGGCGCCTGGT-CACC-3') and reverse (5'-CGCTCCTGGAAGATGGT-GATGG-3').



GMJ Medicine

Figure 1. Effects of different levels of carvacrol on plasma glucose (mg/dL) in diabetic rats

Statistical analysis

The data were analyzed by one-way Analysis of Variance (ANOVA) and post-hoc was conducted by the Dunnett multiple comparison tests by the SPSS statistical package and graphs were illustrated by Graph Pad Software.

Results

Plasma glucose

The data for glucose concentration are presented in Figure 1. Results showed that the plasma concentration of glucose was significantly increased in diabetic Wistar rats in comparison to control group (P<0.05). Oral administration of carvacrol could significantly decrease glucose concentration and the best responses were observed in Wistar rats given with highest levels of carvacrol (P<0.05).

Antioxidant status

Our findings for effects of different levels of carvacrol on antioxidant parameters in diabetic rats are shown in Table 1. As results show, diabetic control rats showed lower levels for antioxidant enzymes and higher levels for MDA in comparison to healthy control (P<0.05). Oral administration of carvacrol could alleviate negative effects of diabetes on antioxidant status, so that we did

Table 1. Effects of different levels of carvacrol on antioxidant parameters in diabetic rats

Parameters	Control	Diabetic	Carvacrol-5	Carvacrol-10	Carvacrol-15	P
SOD (U/mg protein)	15.31±1.32 ^a	9.13±0.95 ^d	11.15±0.98 ^c	12.97±0.58 ^b	14.10±0.63 ^a	****
GPx (U/mg protein)	18.61±0.85 ^a	11.53±0.91 ^d	13.65±0.59 ^c	15.21±0.72 ^b	17.93±0.93 ^a	****
CAT (U/mg protein)	204.21±3.25 ^a	142.51±2.35 ^d	158.32±3.24 ^c	171.25±2.94 ^b	198.56±6.51 ^a	****
MDA (nmol/mg protein)	19.08±1.31 ^c	26.34±1.15 ^a	22.15±1.57 ^b	21.45±2.13 ^b	19.78±0.63 ^c	****

Superscripts (a-d) show significant differences at P<0.05 in per row.

GMJ




Figure 2. Effects of different levels of carvacrol on pro-inflammatory cytokines in diabetic rats

GMJ
Medicine

not observe significant difference between healthy control rats and those received carvacrol in level of 15 mg/kg ($P > 0.05$).

Pro-inflammatory cytokines gene expression

Effects of different levels of carvacrol on pro-inflammatory cytokines in diabetic rats are shown in Figure 2. As results indicate diabetes increased levels of pro-inflammatory cytokines but oral administration of carvacrol especially in higher levels could alleviate effects of diabetes on pro-inflammatory cytokines.

Discussion

Administration of streptozotocin could significantly increase levels of glucose and MDA. MDA is end product for lipid peroxidation. There is a correlation exists between lipid peroxidation and increased glucose. Manana et al. showed increased lipid peroxidation in tissue that attributed it to increased blood glucose [21]. Lipid peroxidation has been known during diabetes mellitus which is one key feature in patients with diabetes mellitus. The free radical and Reactive Oxygen Species modulate in the different types of diseases such as diabetes. The free radicals could have important role in diabetes mellitus [22].

Lipid peroxidation is known to have risks for body. Increased lipid peroxidation in the tissues of diabetic rats could be attributed to increased TBARS and hydroperoxides in the tissues [23]. With regards to increased glucose, it has been reported a positive correlation between hyperglycemia and inflammation [24]. It means that inflammation stimulates insulin resistance in the physiological level [25, 26]. Inflammation is involved in progression of diabetes and increases hyperglycemia. Oral administration of carvacrol could significantly decrease levels of glucose.

Ezhumalai et al. have reported that administration of carvacrol significantly decreased level of glucose and glycosylated hemoglobin [27]. Improved levels of glu-

cose could be attributed to antioxidant status and inflammatory responses. Results showed that inflammatory responses and antioxidant status were significantly improved in Wistar rats fed with higher levels of carvacrol. It has been reported decreased activities of enzymatic antioxidants during diabetes, and it could be attributed to a number of deleterious effects due to the increased free radicals [28].

The SOD and CAT are known as two key enzymes that remove the toxic free radicals induced by STZ. The activities of SOD and CAT were decreased in liver tissue of diabetic rats [29]. SOD is involved in conversion of the superoxide radicals into H_2O_2 and molecular oxygen and also maintains the tissues from highly reactive hydroxyl radical by catalyzing the decreased hydrogen peroxides [29].

In the present study, carvacrol improved glucose level and MDA by improving level of antioxidants. Similarly, Deng et al. reported that the use of carvacrol could prevent increased oxidative stress, also TNF- α and NF- κ B signaling in diabetic rats [30-32]. As mentioned earlier that diabetes is related with inflammation and thus our result shows that carvacrol decreases inflammation in diabetic rat model.

Conclusion

In summary, the carvacrol as novel agent could help animals with diabetes in terms of hyperglycemia, inflammation and oxidation. Since higher levels of carvacrol could significantly show anti-inflammatory and antioxidant properties, we thus recommend it to be included in the commercial preparations for prescription as therapeutic agents.

Ethical Considerations

Compliance with ethical guidelines

There was no ethical considerations to be considered in this research.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

All authors contributed toward data analysis, drafting and revising the paper and agreed to be responsible for all the aspects of this work.

Conflict of interest

The authors declared no conflict of interest.

References

- [1] Kahraman C, Yümün G, Kahraman N, Namdar N, Cosgun S. Neutrophil-tolymphocyteratio in diabetes mellitus patients with and without diabetic footulcer. *Eur J Med Sci*. 2014; 1(1):8-13. [DOI:10.12973/ejms.2014.102p]
- [2] Colhoun H, Schalkwijk C, Rubens M, Stehouwer C. C-reactive protein in type 1 diabetes and its relationship to coronary artery calcification. *Diabetes Care*. 2002; 25:1813-7. [DOI:10.2337/diacare.25.10.1813] [PMID]
- [3] Hayaishi-Okano R, Yamasaki Y, Katakami N, Ohtoshi K, Gorogawa S. Elevated Creactive protein associates with early-stage carotid atherosclerosis in young subjects with type 1 diabetes. *Diabetes Care*. 2002; 25:1432-8. [DOI:10.2337/diacare.25.8.1432] [PMID]
- [4] Jialal I, Devaraj S, Venugopal S. Oxidative stress, inflammation and diabetic vasculopathies: The role of alpha tocopherol therapy. *Free Radic Res*. 2002; 36(12):1331-6. [DOI:10.1080/1071576021000038531] [PMID]
- [5] Ghaderian SB, Hayati F, Shayanpour S, Mousavi SSB. Diabetes and end-stage renal disease: A review article on new concepts. *J Renal Inj Prev*. 2015; 4:28-33.
- [6] Magri CJ, Fava S. The role of tubular injury in diabetic nephropathy. *Eur J Intern Med*. 2009; 20:551-5. [DOI:10.1016/j.ejim.2008.12.012] [PMID]
- [7] Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res*. 2010; 107:1058-70. [DOI:10.1161/CIRCRESA-HA.110.223545] [PMID] [PMCID]
- [8] Wada J, Makino H. Inflammation and the pathogenesis of diabetic nephropathy. *Clin Sci*. 2013; 124:139-52. [DOI:10.1042/CS20120198] [PMID]
- [9] Schmidt AM, Yan SD, Wautier JL, Stern DM. Activation of receptor for advanced glycation end products—a mechanism for chronic vascular dysfunction in diabetic vasculopathy and atherosclerosis. *Circ Res*. 1999; 84:489-97. [DOI:10.1161/01.RES.84.5.489] [PMID]
- [10] Romano M, Pomilio M, Vigneri S, Falco A, Chiesa PL, Chiarelli F, et al. Endothelial perturbation in children and adolescents with type 1 diabetes: Association with markers of the inflammatory reaction. *Diabetes Care*. 2001; 24:1674-8. [DOI:10.2337/diacare.24.9.1674] [PMID]
- [11] Eo H, Lee H-J, Lim Y. Ameliorative effect of dietary genistein on diabetes induced hyper-inflammation and oxidative stress during early stage of woundhealing in alloxan induced diabetic mice. *Biochem Biophys Res Commun*. 2016; 478: 1021-7. [DOI:10.1016/j.bbrc.2016.07.039] [PMID]
- [12] Rahimi R, Nikfar S, Larijani B, Abdollahi M. A review on the role of antioxidants in the management of diabetes and its complications. *Biomed Pharmacotherap*. 2005; 59: 365-73. [DOI:10.1016/j.biopha.2005.07.002] [PMID]
- [13] Amiri R, Akbari M. The role of carvacrol as active compound of essential oils in diabetes. *Biomed J Sci Tech Res*. 2018; 11(1): [DOI:10.26717/BJSTR.2018.11.002054]
- [14] Kirimer N, Baser KHC, Tumen G. Carvacrol rich plants in Turkey. *Chem Nat Comp*. 1995; 31(1):37-42. [DOI:10.1007/BF01167568]
- [15] Berger JP, Akiyama TE, Meinke PT. PPARs: Therapeutic targets for metabolic disease. *Trends Pharmacol Sci*. 2005; 26(5):244-51. [DOI:10.1016/j.tips.2005.03.003] [PMID]
- [16] Kallistratos G, Evangelou A, Agnantis N, Fasske E, Karkabounas S, et al. Enhancement of the antineoplastic effect of anticarcinogens on benzo(a)pyrene-treated Wistar rats, in relation to their number and biological activity. *Cancer Lett*. 1994; 82:153-65. [DOI:10.1016/0304-3835(94)90006-X]
- [17] Evangelou A, Kalpouzos G, Karkabounas S, Liasko R, Nonni A, et al. Dose-related preventive and therapeutic effects of antioxidants-anticarcinogens on experimentally induced malignant tumors in Wistar rats. *Cancer Lett*. 1997; 115:105-11. [DOI:10.1016/S0304-3835(97)04712-5]
- [18] Mansouri E, Panahi M, Ghaffari MA, Ghorbani A. Effects of grape seed proanthocyanidin extract on oxidative stress induced by diabetes in rat kidney. *Iran Biomed J*. 2011; 15:100-6.
- [19] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochem*. 1976; 72:248-54. [DOI:10.1016/0003-2697(76)90527-3]
- [20] Khan HA, Abdelhalim MAK, Alhomida AS, Al Ayed MS. Transient increase in IL-1 β , IL-6 and TNF- α gene expression in rat liver exposed to gold nanoparticles. *Genetic Mol Res*. 2013; 12(4):5851-7. [DOI:10.4238/2013.November.22.12] [PMID]
- [21] Manna P, Ghosh J, Das J, Sil PC. Streptozotocin induced activation of oxidative stress responsive splenic cell signaling pathways: Protective role of arjunolic acid. *Toxicol Appl Pharmacol*. 2010; 244:114-29. [DOI:10.1016/j.taap.2009.12.024] [PMID]
- [22] Yang H, Jin X, Lam CWK, Yan SK. Oxidative stress and diabetes mellitus. *Clin Chem Lab Med*. 2011; 49:1773-82. [DOI:10.1515/cclm.2011.250] [PMID]
- [23] Arulselvan P, Subramanian S. Ultrastructural and biochemical abnormalities in the liver of streptozotocin-diabetic rats: Protective effects of *Murraya koenigii*. *J Pharmacol Toxicol*. 2008; 3:190-202. [DOI:10.3923/jpt.2008.190.202]
- [24] Saxena M, Modi D. Inflammation and diabetes. *Interdiscip J Microinflammation*. 2014; 1:110.
- [25] Wu T, Dorn J, Donahue R, Sempos C, Trevisan M. Associations of serum Creactiveprotein with fasting insulin, glucose, and glycosylat-

- ed hemoglobin. *Am J Epidemiol.* 2002; 155:65-71. [DOI:10.1093/aje/155.1.65] [PMID]
- [26] Fichtlscherer S, Rosenberger G, Walter D, Breuer S, Dimmeler S, Zeiher A. Elevated C-reactive protein levels and impaired endothelial vasoreactivity in patients with coronary artery disease. *Circulation.* 2000; 102:1000-6. [DOI:10.1161/01.CIR.102.9.1000] [PMID]
- [27] Ezhumalai M, Radhiga T, Viswanathan Pugalendi K. Antihyperglycemic effect of carvacrol in combination with rosiglitazone in high-fat diet-induced type 2 diabetic C57BL/6J mice. *Mol Cell Biochem.* 2014; 385:23-31. [DOI:10.1007/s11010-013-1810-8] [PMID]
- [28] Srinivasan S, Pari L. Ameliorative effect of diosmin, acitrus flavonoid against streptozotocin-nicotinamide generated oxidative stress induced diabetic rats. *Chemico-Biol Interact.* 2012; 95:43-51. [DOI:10.1016/j.cbi.2011.10.003] [PMID]
- [29] Omotayo EO, Gurtu S, Sulaiman SA, Wahab MSA, Sirajudeen KNS, Salleh MSM. Hypoglycemic and antioxidant effects of honey supplementation in streptozotocin-induced diabetic rats. *Int J Vitamin Nutr Res.* 2010; 80:74-82. [DOI:10.1024/0300-9831/a000008] [PMID]
- [30] Deng W, Lu H, Teng J. Carvacrol attenuates diabetes-associated cognitive deficits in rats. *J Mol Neurosci.* 2009; 51(3):813-9. [DOI:10.1007/s12031-013-0069-6] [PMID]
- [31] Nasirian F, Dadkhah M, Moradi-Kor N, Obeidavi Z. Effects of *Spirulina platensis* microalgae on antioxidant and anti-inflammatory factors in diabetic rats. *Diabetes Metab Syndr Obes.* 2018; 11:375-80. [DOI:10.2147/DMSO.S172104] [PMID] [PMCID]
- [32] Mesbahzadeh B, Rajaei SA, Tarahomi P, Seyedinia SA, Rahmani M, Rezamohamadi F, et al. Beneficial effects of *spirogyra neglecta* extract on antioxidant and anti-inflammatory factors in streptozotocin-induced diabetic rats. *Biomolecular Concepts.* 2018; 9(1):184-9. [DOI:10.1515/bmc-2018-0015] [PMID]