

Original Article: Comparison of Ondansetron and Metoclopramide in the Prevention of Nausea and Vomiting after Coronary Artery Bypass Graft Surgery

Sahar Sadeghi¹, Hassan Mohammadipour Anvari^{2,*}

¹Anesthesiologist, Tuberculosis and Lung Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Associate Professor of Anesthesiology, Tuberculosis and Lung Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran



Citation S. Sadeghi, H. Mohammadipour Anvari*, Comparison of ondansetron and metoclopramide in the prevention of nausea and vomiting after coronary artery bypass graft surgery. *GMJ Med.* 2021, 5(1):440-444.

<http://doi.org/10.22034/GMIM.2021.1.29>



Article info:

Received: 29 September 2021

Accepted: 16 December 2021

Available Online: 17 December 2021

Checked for Plagiarism: Yes,

Language Editor: Ermia Aghaie

Editor who Approved Publication:

Professor Dr. Ali Nokhodchi

Keywords:

Metoclopramide, Ondansetron, PONV, CABG

ABSTRACT

Introduction: The aim of this study was to determine the effect of ondansetron and metoclopramide on the prevention of PONV in high-risk patients. The reason for choosing these two drugs is that ondansetron is a drug that is widely used in the world today as a first-line drug in the prevention and treatment of PONV, and metoclopramide is a drug that is cheap and available as a common anti-emetic drug by surgeons. PONV control is prescribed in Iran.

Material and Methods: A total of 126 patients who were candidates for coronary artery bypass graft surgery during 2018 and 2019 were evaluated in this study. Intraoperative injection of metoclopramide or ondansetron was compared between all patients and the severity of postoperative nausea and vomiting was compared.

Results: The rate of nausea in patients was about 67.5% with an average of 158.5 seconds of nausea in these patients each day and there was no significant difference between the two drug groups in terms of these two variables (P-Value=0.342 and P-Value=0.900). The average rate of vomiting in patients was 19.8%, of which 52% had vomiting once and 48% had vomited more than once. There was a significant difference in the presence of vomiting in the two drug groups (P-Value=0.044).

Conclusion: The results of this study showed that metoclopramide and ondansetron have similar effects in contrast to nausea and vomiting after coronary artery bypass graft surgery.

Material and Methods

*Corresponding Author: Hassan Mohammadipour Anvari (dr.anvaritbzmed@yahoo.com)

Introduction

One of the major problems for patients and medical staff following surgery is postoperative nausea and vomiting (PONV), and is a major concern for patients about surgery and anesthesia. In general, postoperative nausea and vomiting occur in 20 to 30% of patients who undergo surgery, although there are many differences in the values reported in different studies [1,2].

Today, based on the results of extensive studies, a clear guideline has been published on how to administer PONV, according to which the prophylactic administration of antimycotic drugs is recommended for patients at moderate and high risk. The antimicrobial drugs used inhibit one or more neurotransmitter regions in the chemoreceptor trigger zone in the medulla and include anticholinergics, antipopaminergics, antihistaminergics, and antiserotonergics [3,4].

Because at present in our country there is no specific protocol for how to administer PONV, and considering that the routine method of anesthesia is the use of general anesthesia with inhaled drugs with nitric oxide (NO) as well as opioids as Routine analgesics are used to control pain in surgical patients [5]; patients are at moderate to high risk for PONV (regardless of patient-related risk factors and surgery). As a result, prophylactic treatment for these patients is indicated, according to a guideline published by the American Society of Anesthesiologists [6].

The aim of this study was to determine the effect of ondansetron and metoclopramide on the prevention of PONV in high-risk patients. The reason for choosing these two drugs is that ondansetron is a drug that is widely used in the world today as a first-line drug in the prevention and treatment of PONV, and metoclopramide is a drug that is cheap and available as a common anti-emetic drug by surgeons PONV control is prescribed in Iran.

In this prospective descriptive study, 126 patients in Class 1 and 2 of the American Society of Anesthesiologists (ASA I-II) aged 18 to 65 years underwent coronary artery bypass graft surgery and received opioid medications during surgery. , From 2018 to 2019, are examined in the surgical ward of Shahid Madani Hospital, and patients who have contraindications to the use of ondansetron or metoclopramide, have a history of taking drugs that interact with ondansetron or metoclopramide, a history of allergy to ondansetron or metoclopramide Pregnant and lactating women and patients who have a mechanical cause for postoperative nausea and vomiting (such as premature obstruction or ileus) are excluded from the study. The patients in question are visited by a respected anesthesiologist the day before surgery and are included in the study if they meet the criteria for selecting a research sample. The studied patients are completely randomly divided into 2 equal groups of 63 patients. (Using a table of random numbers by computer)

In the operating room, general anesthesia is induced with a standard dose of inhaled anesthetic and maintained with NO. It is prescribed as an intraoperative analgesic for the opioid patient. The desired anti-nausea drugs (ondansetron 4mg or metoclopramide) 10mg are taken by the anesthesiologist in the required number in equal volumes into the syringe and are prepared according to the patient's number to the anesthesiologist, who does not know the type of drug, and The drug is injected intravenously about 30 minutes before surgery or immediately after induction of anesthesia if the duration of surgery is less than half an hour. The PACU is cared for and then transferred to the general surgery ward. In order to control the confounding variables, all patients should be restricted to receiving food orally for at least 8 hours before to 24 hours after surgery and also be sufficiently hydrated and receive opioid analgesics if pain persists after surgery. Absolute rest for at least 24 hours after surgery. The patients were visited 24 hours after the surgery

by the relevant intern, who did not know the patient's medication, and the patient questionnaire form was based on information obtained from patients and their companions, as well as the nurses' report, which The patient file will be registered and completed. The patients to be admitted to the study were visited by a respected anesthesiologist the day before surgery, and if they met the criteria for selecting the research sample, they were provided with complete and necessary explanations on how to perform the study and then written consent. It was taken from them.

After confirmation of this research in the ethics committee of Tabriz University of Medical Sciences (IR.TBZMED.REC.1399.980), the objectives of the research were explained to patients in simple language. Written informed consent was completed by patients.

In the analysis of qualitative variables from Chi-Square statistical tests (Chi-Square) and Fisher

Exact test (Fisher Exact test) and in the analysis of quantitative variables from t-test and non-parametric alternative tests suitable when needed using Spss and Statistica statistical software were used.

Results

The rate of nausea in patients was about 67.5% with an average of 158.5 seconds of nausea in these patients each day and there was no significant difference between the two drug groups in terms of these two variables (P-Value=0.342 and P-Value=0.900). The average rate of vomiting in patients was 19.8%, of which 52% had vomiting once and 48% had vomited more than once. There was a significant difference in the presence of vomiting in the two drug groups (P-Value=0.044) but no significant difference was observed in the number of vomiting (P-Value=0.097). Details of vomiting and nausea in patients can be seen in Table 2.

Table 1: Comparison of the severity of nausea and vomiting between study participants

Drug	Total		ondansetron		metoclopramide	
nausea and vomiting	%	N	%	N	%	N
No	65.9%	83	71.4%	45	60.3%	38
nausea and vomiting	18.3%	23	12.7%	8	23.8%	15
Just nausea	14.3%	18	15.9%	10	12.7%	8
Just vomiting	1.6%	2	0	0	3.2%	2
Total	100%	126	100%	63	100%	63
P Value	Fisher's Exact Test=4.54 P-Value=0.194					

Discussion

The etiology of postoperative nausea and vomiting following open abdominal surgery depends on several factors, including patient-related factors, factors related to anesthesia technique, factors related to the type of surgery, and factors after surgery [7]. Because at present in our country there is no specific protocol for how to manage nausea and vomiting after surgery, and considering that the routine method of anesthesia is the use of general anesthesia with inhaled drugs and also opioid drugs as a routine analgesic Used to control pain

in surgical patients, patients are at moderate to high risk for nausea and vomiting after surgery (regardless of patient-related risk factors and surgery) [8]. As a result, prophylactic treatment for these patients is indicated, according to a guideline published by the American Society of Anesthesiologists [9,10]. Therefore, we decided to compare the effects of ondansetron and metoclopramide in the prevention of postoperative nausea and vomiting in a present study [11,12]. The present study is a prospective study that investigated the effect of ondansetron and metoclopramide on the prevention of nausea and vomiting after coronary artery

bypass graft surgery. In this study, 126 patients in two groups of 63 were studied [13].

The results of this study showed that there was no significant difference between the two drug groups in terms of nausea in the first 24 hours after open abdominal surgery in patients receiving ondansetron 30 minutes before the end of surgery compared to metoclopramide. However, the incidence of vomiting during this period was lower in patients receiving ondansetron than in metoclopramide, and there was a significant difference between the two drug groups in terms of vomiting. In general, there is no significant difference between the two drug groups in terms of nausea and vomiting [14,15]. In a recent study, both groups were similar in terms of risk factors for nausea and vomiting after surgery and there was a significant difference between the two drug groups in terms of most patient-related risk factors (sex, history of motion sickness, history of chronic nausea and vomiting, history). Nausea and vomiting in previous surgeries and smoking) and the type of surgery is not observed. In addition, the technique of induction and maintenance of anesthesia and the method of analgesic use during surgery were the same for all patients and all patients had the same conditions in terms of postoperative factors. As a result, the difference in the incidence of postoperative nausea and vomiting is mainly due to prophylactic antiemetic drugs prescribed during surgery [1,16].

Because the side effects of ondansetron and metoclopramide are short-lived and self-limiting, the side effects of the prescribed anti-nausea drugs were not evaluated in this study. In general, the results of our study show that the two drugs ondansetron at a dose of 4 mg and metoclopramide at a dose of 10 mg in case of intravenous injection 30 minutes before surgery do not have a superior advantage over each other in preventing nausea and vomiting after surgery. Because metoclopramide is more economical than ondansetron, its use is recommended to prevent postoperative nausea and vomiting in high-risk patients [12,14]. It is suggested that other studies with a larger sample size be performed in the future to confirm the accuracy of the results obtained

from this study. In addition, in these trials, it is better to carefully control all the factors that may affect the outcome of the work (type of surgery, anesthesia technique and facto-release after surgery) and to be extremely careful in selecting patients. Another important point is the dose and time of injection of anti-nausea drugs, which should be adjusted according to the half-life and peak time of the drug.

Conclusion

The results of this study showed that metoclopramide and ondansetron have similar effects in contrast to nausea and vomiting after coronary artery bypass graft surgery.

References

- [1] S.R. Pitts, *Ann. Intern. Med.*, **2014**, *161*, JC3 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [2] M. Talebpour, N. Ghasnejad Omrani, F. Imani, R. Shariat Moharari, P. Pourfakhr, M.R. Khajavi, *Anesth. Pain. Med.*, **2017**, *7*, e57810 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [3] S. Maitra, A. Som, D.K. Baidya, S. Bhattacharjee, *Anesthesiol. Res. Pract.*, **2016**, *2016*, 7089454 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [4] S. Dalhat, A. Mohammad, *Niger. J. Basic Clin. Sci.*, **2018**, *15*, 24-28 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5] A. Reza, E. Riahi, A. Daneshi, E. Golchini, *Brain Inj.*, **2018**, *32*, 487-492 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6] B. Balakrishnan, R.M. Rus, K.H. Chan, A.G. Martin, M.S. Awang, *Asian J. Neurosurg.*, **2019**, *14*, 201-205 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7] S. Bagheri-Hariri, M. Bahreini, P. Farshidmehr, S. Barzandeh, S. Babaniamansour, E. Aliniagerdroudbari, A. Baratloo, *Arch. Trauma Res.*, **2019**, *8*, 207-213 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8] E. Farzaneh, G. Fattahzadeh-Ardalani, V. Abbasi, F. Kahnamouei-aghdam, B. Molaei, E. Iziy, H. Ojaghi, *Emerg. Med. Int.*, **2017**, *2017*, 1439486 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [9] H. Gerritsen, M. Samim, H. Peters, H. Schers, F.A. van de Laar, *BMJ Open*, **2018**, *8*, e020364 [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [10] M. Zamani, B. Namdar, R. Azizkhani, O. Ahmadi, M. Esmailian, *Emergency*, **2015**, *3*, 137-140 [crossref], [Google Scholar], [Publisher]
- [15] P. Tienchaiananda, W. Nipondhkit, K. Maneenil, S. Sa-Nguansai, S. Payapwattanawong, *Ann. Palliat. Med.*, **2019**, *8*, 372-380 [crossref], [Google Scholar], [Publisher]

Copyright © 2021 by SPC ([Sami Publishing Company](#)) + is an open access article distributed under the Creative Commons Attribution License(CC BY) license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

- [11] H. Hashimoto, M. Abe, O. Tokyama, H. Mizutani, Y. Uchitomi, T. Yamaguchi, Y. Hoshina, Y. Sakata, T.Y. Takahashi, K. Nakashima, M. Nakao, *Lancet Oncol.*, **2020**, *21*, 242-249 [crossref], [Google Scholar], [Publisher]
- [12] S. Ithimakin, P. Theeratrakul, A. Laocharoenkiat, A. Nimmannit, C. Akewanlop, N. Soparattanapaisarn, S. Techawattanawanna, K. Korphaisarn, P. Danchaivijitr, *Support Care Cancer*, **2020**, *28*, 5335-5342 [crossref], [Google Scholar], [Publisher]
- [13] S.Y. Jeon, H.S. Han, W.K. Bae, M.R. Park, H. Shim, S.C. Lee, S.I. Go, H.J. Yun, Y.J. Im, E.K. Song, *Cancer Res. Treat.*, **2019**, *51*, 90-97 [crossref], [Google Scholar], [Publisher]
- [14] S. Mukhopadhyay, K.P.A. Kwatra, D. Badyal, *Support Care Cancer*, **2017**, *25*, 145-154 [crossref], [Google Scholar], [Publisher]
- [16] R.M. Navari, C.M. Pywell, J.G. Le-Rademacher, P. White, A.B. Dodge, C. Albany, C.L. Loprinzi, *JAMA. Oncol.*, **2020**, *6*, 895-899 [crossref], [Google Scholar], [Publisher]
- [17] N. Kalakonda, M. Maerevoet, F. Cavallo, G. Follows, A. Goy, J.S. Vermaat, O. Casasnovas, N. Hamad, J.M. Zijlstra, S. Bakhshi, R. Bouabdallah, *Lancet Haematol.*, **2020**, *7*, e511-e522 [crossref], [Google Scholar], [Publisher]
- [18] A. Chari, D.T. Vogl, M. Gavriatopoulou, A.K. Nooka, A.J. Yee, C.A. Huff, P. Moreau, D. Dingli, C. Cole, S. Lonial, M. Dimopoulos, *N. Engl. J. Med.*, **2019**, *381*, 727-738 [crossref], [Google Scholar], [Publisher]